

Differential inequality of the second derivative that leads to normality

Qiaoyu Chen, Shahar Nevo, Xuecheng Pang

Abstract

Let \mathcal{F} be a family of functions meromorphic in a domain D . If $\{\frac{|f''|}{1+|f|^3} : f \in \mathcal{F}\}$ is locally uniformly bounded away from zero, then \mathcal{F} is normal.

I. Introduction.

Recently, progress was occurred concerning the study of the connection between differential inequalities and normality. A natural point of departure for this subject is the well-known theorem due to F.Marty.

Marty's Theorem [8, P.75] A family \mathcal{F} of functions meromorphic in a domain D is normal if and only if $\{f^\# : f \in \mathcal{F}\}$ is locally uniformly bounded in D .

Following Marty's Theorem, L. Royden proved the following generalization.

Theorem R[7] Let \mathcal{F} be a family of functions meromorphic in a domain D , with the property that for each compact set $K \subset D$, there is a positive increasing function h_K , such that $|f'(z)| \leq h_K(|f(z)|)$ for all $f \in \mathcal{F}$ and $z \in K$. Then \mathcal{F} is normal in D .

This result was significantly extended further in various directions, see [3], [9] and [11]. S.Y.Li and H.Xie established a different kind of generalization of Marty's Theorem that involves higher derivatives.

2010 Mathematical subject classification 30A10, 30D45.

keywords and phrases, Normal family, Differential inequality.

Theorem LX [4] Let \mathcal{F} be a family of functions meromorphic in a domain D , such that each $f \in \mathcal{F}$ has zeros only of multiplicities $\geq k$, $k \in \mathbb{N}$. Then \mathcal{F} is normal in D if and only if the family

$$\left\{ \frac{|f^{(k)}(z)|}{1 + |f(z)|^{k+1}} : f \in \mathcal{F} \right\}$$

is locally uniformly bounded in D .

In [6], the second and the third authors gave a counterexample to the validity of Theorem LX, without the condition on the multiplicities of zeros for the case $k = 2$.

Concerning differential inequalities with the reversed sign of the inequality, J. Grahl, and the second author proved the following result, that may be considered as a counterpart to Marty's Theorem.

Theorem GN [1] Let \mathcal{F} be a family of functions meromorphic in D , and $c > 0$. If $f^\#(z) > c$ for every $f \in \mathcal{F}$ and $z \in D$, then \mathcal{F} is normal in D .

N. Steinmetz [10], gave a shorter proof of Theorem GN, using the Schwarzian derivative and some well-known facts on linear differential equations.

Then in [5], X.J. Liu together with the second and third authors generalized Theorem GN and proved the following result.

Theorem LNP Let $1 \leq \alpha < \infty$ and $c > 0$. Let \mathcal{F} be the family of all meromorphic functions f in D , such that

$$\frac{|f'(z)|}{1 + |f(z)|^\alpha} > c$$

for every $z \in D$.

Then the following hold:

- (1) If $\alpha > 1$, then \mathcal{F} is normal in D .
- (2) If $\alpha = 1$, then \mathcal{F} is quasi-normal in D but not necessarily normal.

Observe that (2) of the theorem is a differential inequality that distinguishes between quasi-normality and normality.

In this paper, we continue to study differential inequality with the reversed sign (" \geq ") and prove the following theorem.

Theorem 1. Let D be a domain in \mathbb{C} and let $c > 0$. Then the family \mathcal{F} of all functions f

meromorphic in D , such that

$$\frac{|f''(z)|}{1 + |f(z)|^3} > C$$

for every $z \in D$ is normal.

Observe that the above differential inequality is the reversed inequality to that of Theorem LX in the case $k = 2$.

Let us set some notation.

For $z_0 \in C$ and $r > 0$. $\Delta(z_0, r) = \{z : |z - z_0| < r\}$, $\overline{\Delta}(z_0, r) = \{z : |z - z_0| \leq r\}$. We write $f_n(z) \xrightarrow{X} f(z)$ on D to indicate that the sequence $\{f_n(z)\}$ converges to $f(z)$ in the spherical metric, uniformly on compact subsets of D , and $f_n(z) \Rightarrow f(z)$ on D if the convergence is also in the Euclidean metric.

II Proof of Theorem 1.

Since $|f''| > c$ for every $f \in \mathcal{F}$, it follows that $\{f'' : f \in \mathcal{F}\}$ is normal in D . Let $\{f_n\}_{n=1}^\infty$ be a sequence of functions from \mathcal{F} . Without loss of generality, we can assume that $f_n''(z) \xrightarrow{X} H$ in D .

Let us separate into two cases.

Case 1. $f_n, n \geq 1$ are holomorphic functions in D .

Case 1.1 H is holomorphic function in D .

Since normality is a local property. It is enough to prove that $\{f_n\}$ is normal at each point of D . Let $z_0 \in D$ without loss of generality, we can assume that $z_0 = 0$. By the assumption on H , there exist some $r > 0$, $M > C$, such that $|f_n''(z)| \leq M$ for every $z \in \Delta(0, r)$ if n is large enough. We then get for large enough n and $z \in \Delta(0, r)$ that $1 + |f_n(z)|^3 \leq \frac{2M}{C}$ and we deduce that $\{f_n\}_{n=1}^\infty$ is normal at $z = 0$, as required.

Case 1.2 $H \equiv \infty$ in D .

Again, let $z_0 \in D$ and assume that $z_0 = 0$. Let $r > 0$ be such that $\overline{\Delta}(0, r) \subset D$. Without loss of generality, we can assume that $|f_n''(z)| > 1$ for every $z \in \Delta(0, r)$, $n \in \mathbb{N}$. Then $\log |f_n''|$ is a positive harmonic function in $\Delta(0, r)$.

From Harnack's inequality we then get that

(1)

$$|f_n''(z)| \leq |f_n''(0)|^{\frac{1+|z|}{1-|z|}}$$

for every $z \in \Delta(0, r)$, $n \in \mathbb{N}$.

Let us fix some $0 < \rho < \frac{r}{2}$. Then

(2)

$$\frac{r + \rho}{r - \rho} < 3.$$

For every $n \geq 1$, let $z_n \in \{z : |z| = \rho\}$ be such that

$$|f_n(z_n)| = \max_{|z| \leq \rho} |f_n(z)| = M(\rho, f_n)$$

By Cauchy's Inequality, we get that

$$|f_n''(0)| \leq \frac{2}{\rho^2} M(\rho, f_n) = \frac{2}{\rho^2} |f_n(z_n)|.$$

Hence, by (1), we get

$$C \leq \frac{|f_n''(z_n)|}{1 + |f_n(z_n)|^3} \leq \frac{|f_n''(z_n)|}{|f_n(z_n)|^3} \leq \frac{|f_n''(0)|^{\frac{r+\rho}{r-\rho}}}{|f_n(z_n)|^3} \leq \left(\frac{2}{\rho^2}\right)^{\frac{r+\rho}{r-\rho}} |f_n(z_n)|^{\frac{r+\rho}{r-\rho}-3},$$

Thus, by (2)

$$M(\rho, f_n) = |f_n(z_n)| \leq \left(\frac{1}{C} \left(\frac{2}{\rho^2}\right)^{\frac{r+\rho}{r-\rho}}\right)^{\frac{1}{3 - \frac{r+\rho}{r-\rho}}},$$

which means that $\{f_n\}$ is locally uniformly bounded in $\Delta(0, \rho)$ and thus $\{f_n\}$ is normal at $z = 0$.

Case 2 f_n are meromorphic functions with pole in D .

By Case 1 we have to prove normality only at point z_0 , where $H(z_0) = \infty$. Such points exist if H is a meromorphic function with poles in D or if $H \equiv \infty$. So let z_0 be such that $H(z_0) \equiv \infty$. Without loss of generality, we can assume that $z_0 = 0$. After moving to a subsequence, that without loss of generality will also be denoted by $\{f_n\}_1^\infty$, we can assume that there is a sequence $\zeta_n \rightarrow 0$ such

that $f_n(\zeta_n) = \infty$. For if it was not the case, then for some $\delta > 0$ and large enough n , f_n would be holomorphic in $\Delta(0, \delta)$, and then we would get the asserted normality by case (1).

Also we can assume the existence of

(3) a sequence $\eta_n \rightarrow 0$ such that $f_n(\eta_n) = 0$.

Indeed, since $H(z_0) = \infty$ there exists some $\delta > 0$ such that for large enough n $\min_{z \in \Delta(0, \delta)} |f_n''| > 1$.

Combining it with $f_n \neq 0$ in some neighbourhood of $z = 0$ gives the normality at $z = 0$ by Gu's Criterion [2].

We can also assume that $\{f_n'\}$ is not normal at $z = 0$. Indeed, if $\{f_n'\}$ would be normal at $z = 0$, then by Marty's theorem there exist $r_1 > 0$ and $M > 0$ such that for large enough n , $\frac{|f_n''(z)|}{1 + |f_n'(z)|^2} < M$ for $z \in \Delta(0, r_1)$. Since $H(0) = \infty$, there exists some $r_2 \leq r_1$ such that for large enough n , $|f_n''(z)| \geq 2M$ for $z \in \Delta(0, r_2)$.

We thus have for large enough n and $z \in \Delta(0, r_2)$, $1 + |f_n'(z)|^2 > \frac{|f_n''(z)|}{M} \geq 2$ and thus $|f_n'(z)| \geq 1$. We then get

$$\frac{|f_n'(z)|^2}{|f_n''(z)|} = \frac{|f_n'(z)|^2}{1 + |f_n'(z)|^2} \cdot \frac{1 + |f_n'(z)|^2}{|f_n''(z)|} \geq \frac{1^2}{1 + 1^2} \cdot \frac{1}{M} = \frac{1}{2M}.$$

Hence We have for large enough n and $z \in \Delta(0, r_2)$

$$(4) \quad \frac{|f_n'(z)|^2}{1 + |f_n(z)|^3} = \frac{|f_n'(z)|^2}{|f_n''(z)|} \cdot \frac{|f_n''(z)|}{1 + |f_n(z)|^3} > \frac{1}{2M} \cdot C.$$

Now, for every $x \geq 0$, $\frac{\sqrt{1+x^2}}{1+x} \geq \frac{1}{\sqrt{2}}$, and by taking square root of (4), we get

$$\frac{|f_n'(z)|}{1 + |f_n(z)|^{\frac{3}{2}}} = \frac{|f_n'(z)|}{\sqrt{1 + |f_n(z)|^3}} \cdot \frac{\sqrt{1 + |f_n(z)|^3}}{1 + |f_n(z)|^{\frac{3}{2}}} > \sqrt{\frac{C}{2M}} \cdot \frac{1}{\sqrt{2}}.$$

By (1) of Theorem LNP, with $\alpha = \frac{3}{2} > 1$, we deduce that $\{f_n\}$ is normal in $\Delta(0, r_2)$ and we are done.

Thus we can assume that $\{f_n'\}$ is not normal at $z = 0$.

Similarly to (3) we can assume that there is a sequence $s_n \rightarrow 0$ such that $f_n'(s_n) = 0$.

We claim that we can assume that $\{\frac{f_n'}{f_n''}\}_{n=1}^\infty$ is not normal at $z = 0$.

Otherwise, after moving to a subsequence that will also be denoted by $\{\frac{f'_n}{f''_n}\}_{n=1}^\infty$ we have $\frac{f'_n}{f''_n} \Rightarrow H_1$ in $\Delta(0, r)$, for some $r > 0$. Since $f''_n \neq 0$ and $\frac{f'_n}{f''_n}(\zeta_n) = 0$ then H_1 must be holomorphic function in $\Delta(0, r)$. Differentiation then gives

$$(5) \quad 1 - \frac{f'_n f''_n}{(f''_n)^2} \Rightarrow H'_1 \text{ in } \Delta(0, r).$$

At $z = s_n$ the left hand of (5) is equal to 1. on the other hand in some small neighbourhood of $z = \zeta_n$, We have $f_n(z) = \frac{A_n}{z - \zeta_n} + \hat{f}_n(z)$, where $A_n \neq 0$ is a constant, and $\hat{f}_n(z)$ is analytic. Here we used that according to the assumption of Theorem 1, all poles of f_n must be simple.

Hence we have $f'_n(z) = \frac{-A_n}{(z - \zeta_n)^2} + \hat{f}'_n(z)$, $f''_n(z) = \frac{2A_n}{(z - \zeta_n)^3} + \hat{f}''_n(z)$, $f_n^{(3)}(z) = \frac{-6A_n}{(z - \zeta_n)^4} + \hat{f}_n^{(3)}(z)$. Then the left hand of (5) get at $z = \zeta_n$. The value $1 - \frac{6}{4} = -\frac{1}{2} \neq 1$, a contradiction.

Claim there exist $r > 0$ and $k > 0$ such that for large enough n , $|\frac{f'_n}{f''_n}(z)|, \left| \frac{f_n^2}{f''_n}(z) \right| \leq K$ for $z \in \Delta(0, r)$.

Proof of Claim Since $H(0) = \infty$, there exist $r > 0$ and $M > 0$ such that $\overline{\Delta}(0, r) \subset D$ and such that for large enough n , $|f''_n(z)| > M$ for $z \in \Delta(0, r)$.

Now, if $|f_n(z)| \leq |f''_n(z)|^{\frac{1}{3}}$ then

$$(6) \quad \left| \frac{f'_n}{f''_n}(z) \right| \leq \frac{|f''_n(z)|^{\frac{1}{3}}}{|f''_n(z)|} \leq \frac{1}{M^{\frac{2}{3}}}$$

and

$$(7) \quad \left| \frac{f_n^2}{f''_n}(z) \right| \leq \frac{|f''_n(z)|^{\frac{2}{3}}}{|f''_n(z)|} \leq \frac{1}{M^{\frac{1}{3}}}.$$

If on the other hand $|f_n(z)| \geq |f''_n(z)|^{\frac{1}{3}}$, then since $\frac{x}{1+x^3} \leq \frac{2^{\frac{2}{3}}}{3}$ for $x \geq 0$, we get

$$(8) \quad \left| \frac{f'_n}{f''_n}(z) \right| = \frac{1 + |f_n(z)|^3}{|f''_n(z)|} \cdot \frac{|f_n(z)|}{1 + |f_n(z)|^3} \leq \frac{1}{C} \cdot \frac{2^{\frac{2}{3}}}{3}.$$

Also We have $\frac{x^2}{1+x^3} \leq \frac{2^{\frac{2}{3}}}{3}$ for $x \geq 0$ and thus

$$(9) \quad \left| \frac{f_n^2}{f''_n}(z) \right| = \frac{1 + |f_n(z)|^3}{|f''_n(z)|} \cdot \frac{|f_n^2(z)|}{1 + |f_n(z)|^3} \leq \frac{1}{C} \cdot \frac{2^{\frac{2}{3}}}{3}.$$

The claim then follows by taking $k = \max \left\{ \frac{1}{M^{\frac{2}{3}}}, \frac{1}{M^{\frac{1}{3}}}, \frac{1}{C} \cdot \frac{2^{\frac{2}{3}}}{3} \right\}$ and consider (6),(7),(8)and (9).

From the claim we deduce that $\{\frac{f'_n}{f''_n}\}_{n=1}^\infty$ and $\{\frac{f_n^2}{f''_n}\}_{n=1}^\infty$ are normal in $\Delta(0, r)$, so after moving to a subsequence, that also will be denote by $\{f_n\}_{n=1}^\infty$, we get that

$$(10) \quad \frac{f_n}{f_n''} \rightarrow H_1 \text{ in } \Delta(0, r)$$

and

$$(11) \quad \frac{f_n^2}{f_n''} \rightarrow H_2 \text{ in } \Delta(0, r)$$

From the claim it follows that H_1 and H_2 are holomorphic in $\Delta(0, r)$.

Differentiating (10) and (11) gives respectively

$$(12) \quad \frac{f_n'}{f_n''} - \frac{f_n^{(3)}}{f_n''} \cdot f_n \Rightarrow H_1' \text{ in } \Delta(0, r)$$

and

$$(13) \quad 2f_n \cdot \frac{f_n'}{f_n''} - f_n^2 \cdot \frac{f_n^{(3)}}{f_n''^2} \Rightarrow H_2' \text{ in } \Delta(0, r).$$

Since $\{f_n''\}_{n=1}^\infty$ is normal, there exists some $k_1 > 0$ such that $\frac{|f_n^{(3)}(z)|}{1+|f_n''(z)|} \leq k_1$ for every $n \geq 1$ and for every $z \in \Delta(0, r)$. Since in addition for large enough n , $|f_n''(z)| > M$, then

$$\begin{aligned} \frac{|f_n^{(3)}(z)|}{|f_n''(z)|^2} &= \frac{|f_n^{(3)}(z)|}{1+|f_n''(z)|^2} \frac{1+|f_n''(z)|^2}{|f_n''(z)|^2} \\ &\leq k_1 \left(1 + \frac{1}{M^2}\right) := k_2. \end{aligned}$$

Thus

$$(14) \quad \frac{|f_n''(z)|^2}{|f_n^{(3)}(z)|} \geq \frac{1}{k^2} \text{ for large enough } n.$$

Now since we assume that $\{\frac{f_n'}{f_n''}\}$ is not normal at $z = 0$, then after moving to a subsequence, that also will be denoted by $\{f_n\}_{n=1}^\infty$, we get that there exists a sequence of points $t_n \rightarrow 0$, such that

$$\frac{f_n'}{f_n''}(t_n) := M_n \rightarrow \infty, \quad M_n \in \mathbb{C}.$$

Substituting $z = t_n$ in (12) gives

$$(15) \quad M_n - \frac{f_n^{(3)} \cdot f_n}{f_n''^2}(t_n) := \varepsilon_n \rightarrow H_1'(0).$$

Hence

$$f_n(t_n) = (M_n - \varepsilon_n) \frac{f_n''^2}{f_n^{(3)}}(t_n)$$

From (15) we get, by substituting $z = t_n$ in (13)

$$2(M_n - \varepsilon_n) \frac{f_n''^2}{f_n^{(3)}}(t_n) M_n - (M_n - \varepsilon_n)^2 \left(\frac{f_n''^2}{f_n^{(3)}}(t_n) \right)^2 \frac{f_n^{(3)}}{f_n''^2}(t_n) := \delta_n \rightarrow H_2'(0).$$

From this we get after simplifying

$$(M_n^2 - \varepsilon_n^2) \frac{f_n''^2}{f_n^{(3)}}(t_n) = \delta_n.$$

But by (14) the left hand above tends to ∞ as $n \rightarrow \infty$, while the right hand is bounded, a contradiction.

This completes the proof of Theorem 1.

References

- [1] J. Grahl and S. Nevo, *Spherical derivatives and normal families*, to appear in J. Anal. Math., arXiv:1010.4654
- [2] Y. Gu, *A criterion for normality of families of meromorphic functions*, Sci. Sinica (special issue) **1** (1979), 267-274
- [3] A. Hinkkanen, *Normal families and Ahlfors's Five Island Theorem*, New Zealand J. Math. **22** (1993), 39-41
- [4] S.Y. Li and H. Xie, *On normal families of meromorphic functions*, Acta Math. Sin. **4** (1986), 468-476
- [5] X.J. Liu, S. Nevo and X.C. Pang, *Differential inequalities, normality and quasi-normality*, arXiv: 1111.0844
- [6] S. Nevo and X.C. Pang, *A non explicit counterexample to a problem of quasi-normality*, arXiv: 1111.0841

- [7] H.L. Royden, *A criterion for the normality of a family of meromorphic functions*, Ann. Acad. Sci. Fenn. Ser. A I **10** (1985), 499-500
- [8] J. Schiff, *Normal families*, Springer, New York, 1993
- [9] W. Schwick, *On a normality criterion of H.L. Royden*, New Zealand J. Math **23** (1994), 91-92
- [10] N. Steinmetz, *Normal families and linear differential equations*, to appear in J. Anal. Math.
- [11] L. Zalcman, *Normal families: new perspectives*, Bull. Amer. Math. Soc. (N.S.) **35** (1998), 215-230

QIAOYU CHEN, DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, SHANG HAI 200241, P.R. CHINA

E-mail address: goodluckqiaoyu@126.com

SHAHAR NEVO, DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, 52900 RAMAT-GAN, ISRAEL

E-mail address: nevosh@macs.biu.ac.il

XUECHENG PANG, DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, SHANG HAI 200241, P.R. CHINA

E-mail address: xcpang@math.ecnu.cn